

HRSA's Ryan White HIV/AIDS Program


Female Clients: Ryan White HIV/AIDS Program, 2018

Population Fact Sheet | January 2020

The Health Resources and Services Administration's Ryan White HIV/AIDS Program (RWHAP) provides a comprehensive system of HIV primary medical care, medications, and essential support services for low-income people with HIV. More than half the people with diagnosed HIV in the United States—approximately 519,000 people in 2018—receive services through RWHAP each year. The RWHAP funds grants to states, cities/counties, and local community-based organizations to provide care and treatment services to people with HIV to improve health outcomes and reduce HIV transmission among hard-to-reach populations.



Ryan White HIV/AIDS Program Fast Facts: Female Clients



Females comprise a substantial proportion of RWHAP clients. Of the more than half a million clients served by RWHAP, 26.5 percent are female.

More details about this RWHAP client population are outlined below:

■ The majority of female clients served by RWHAP are from racial/ethnic minority populations. The data show that 84.0 percent of female clients are from racial/ethnic minority populations. 62.1 percent of female clients identify as black/African American, which is higher than the national RWHAP average (47.1 percent), and 19.0 percent of female clients identify as Hispanic/Latino, which is lower than the national RWHAP average (23.2 percent).

- The majority of female clients served by RWHAP are low income. Among female clients served, 70.5 percent are living at or below 100 percent of the federal poverty level, which is higher than the national RWHAP average (61.3 percent).
- The data show that 4.2 percent of female RWHAP clients have unstable housing. This is slightly lower than the national RWHAP average (5.3 percent).
- The RWHAP female client population is aging. Among female RWHAP clients served, 48.2 percent are aged 50 years and older, whereas only 3.6 percent of female RWHAP clients are aged 13–24 years.

Medical care and treatment improve health outcomes and decrease the risk of HIV transmission. People with HIV who take HIV medication daily as prescribed and reach and maintain an undetectable viral load have effectively no risk of sexually transmitting the virus to an HIV-negative partner. In 2018, approximately 86.8 percent of female clients receiving RWHAP HIV medical care are virally suppressed,* which is slightly lower than the national RWHAP average (87.1 percent).

^{*}Viral suppression is defined as a viral load result of less than 200 copies/mL at the most recent test, among people with HIV who had at least one outpatient ambulatory health services visit and one viral load test during the measurement year.