



HIV/AIDS Bureau Performance Measures



CD4 Cell Count performance measure moved to the “Archived” category. (January 2015)

Viral Load Monitoring performance measure moved to the “Archived” category. (May 2015)

Performance Measure:	HIV Drug Resistance Testing Before Initiation of Therapy	National Quality Forum #: None
Percentage of patients, regardless of age, with a diagnosis of HIV who had an HIV drug resistance test performed ¹ before initiation ² of HIV antiretroviral therapy if therapy started during the measurement year		
Numerator:	Number of patients who had an HIV drug resistance test performed at any time before initiation of HIV antiretroviral therapy	
Denominator:	Number of patients, regardless of age, with a diagnosis of HIV who <ul style="list-style-type: none"> • were prescribed HIV antiretroviral therapy during the measurement year for the first time; and • had a medical visit with a provider with prescribing privileges³ at least once in the measurement year 	
Patient Exclusions:	None	
Data Elements:	1. Does the patient, regardless of age, have a diagnosis of HIV/AIDS? (Y/N) <ol style="list-style-type: none"> a. If yes, was the patient seen by a provider with prescribing privileges during the measurement year? (Y/N) <ol style="list-style-type: none"> i. If yes, was HIV antiretroviral therapy prescribed during the measurement year for the first time? (Y/N) <ol style="list-style-type: none"> 1. If yes, was an HIV drug resistance test performed at any time prior to prescribing ARV therapy? (Y/N) <ol style="list-style-type: none"> a. If yes, list date. 	
Comparison Data:	National HIVQUAL: Percentage of patients on ART for a minimum of 12 weeks with one visit in each six month period of the review period who are considered suppressed as derived from the last recorded viral load of the review period; suppressed defined as the viral load is <200 copies/mm ³ (either detectable or undetectable) OR <400 copies/mm ³ (and undetectable) (https://www.ehivqual.org/ and http://nationalqualitycenter.org/). The National HIVQUAL reported the 2011 median as 50%.	
U.S. Department of Health & Human Services Guidelines:	<p><u>Adult guidelines:</u>⁴ “HIV drug-resistance testing is recommended in persons with HIV infection at entry into care regardless of whether antiretroviral therapy (ART) will be initiated immediately or deferred (AII). If therapy is deferred, repeat testing should be considered at the time of ART initiation (CIII).</p> <p>“Genotypic testing is recommended as the preferred resistance testing to guide therapy in antiretroviral (ARV)-naïve patients (AIII).”⁴</p> <p><u>Pediatric guidelines:</u>⁵ “Entry into Care—Baseline Evaluation...Genotype resistance testing should be performed, even if cART is not initiated immediately. Testing at the Time of Switching cART...If regimen is switched because of cART failure (see Recognizing and Managing Antiretroviral Treatment Failure in Management of Children Receiving</p>	



HIV/AIDS Bureau Performance Measures



	Antiretroviral Therapy) resistance testing should be performed while a patient is still receiving the failing regimen to optimize the chance of identifying resistance mutations because resistant strains may revert to wild type within a few weeks of stopping ARV drugs (see Antiretroviral Drug-Resistance Testing).”
Use in Other Federal Programs:	None
References/ Notes:	<p>¹HIV drug resistance testing may occur either during or prior to the measurement year, as long as it is performed before ARV therapy is initiated.</p> <p>²The focus of the measure is on initiation of first antiretroviral regimen for HIV treatment, not prophylaxis or re-initiation.</p> <p>³A “provider with prescribing privileges” is a health care professional who is certified in his/her jurisdiction to prescribe medications.</p> <p>⁴Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the use of antiretroviral agents in HIV-1-infected adults and adolescents. Department of Health and Human Services. Available at http://aidsinfo.nih.gov/ContentFiles/AdultandAdolescentGL.pdf . Section accessed October 14, 2014. C-9.</p> <p>⁵Panel on Antiretroviral Therapy and Medical Management of HIV-Infected Children. Guidelines for the Use of Antiretroviral Agents in Pediatric HIV Infection. Available at http://aidsinfo.nih.gov/contentfiles/lvguidelines/pediatricguidelines.pdf. Section accessed January 26, 2015. D-4 and D-6.</p>



HIV/AIDS Bureau Performance Measures



Performance Measure:	Influenza Immunization ¹	National Quality Forum #: 41
Percentage of patients aged 6 months and older seen for a visit between October 1 and March 31 who received an influenza immunization OR who reported previous receipt of an influenza immunization		
Numerator:	Patients who received an influenza immunization OR who reported previous receipt* of an influenza immunization during the current season *Previous receipt can include: previous receipt of the current season's influenza immunization from another provider OR from same provider prior to the visit to which the measure is applied (typically, prior vaccination would include influenza vaccine given since August 1st).	
Denominator:	All patients aged 6 months and older seen for a visit between October 1 and March 31	
Patient Exclusions:	1. Documentation of medical reason(s) for not receiving influenza immunization (eg, patient allergy, other medical reasons) 2. Documentation of patient reason(s) for not receiving influenza immunization (eg, patient declined, other patient reasons) 3. Documentation of system reason(s) for not receiving influenza immunization (eg, vaccine not available, other system reasons)	
Data Elements:	1. Did the patient, aged six months and older, have at least one medical visit between October 1 and March 31? (Y/N) <ul style="list-style-type: none"> a. Did the patient receive an influenza vaccination or report previous receipt of an influenza vaccination? (Y/N) ***Greater measure specification detail is available including data elements for each value set at https://www.cms.gov/regulations-and-guidance/legislation/ehrincentiveprograms/ecqm_library.html	
Comparison Data:	National HIVQUAL: Percentage of patients who received an influenza vaccination during the review period (https://www.ehivqual.org/ and http://nationalqualitycenter.org/). The National HIVQUAL reported the median as 62% in 2009 and 67% in 2011.	
U.S. Department of Health & Human Services Guidelines:	"Annual vaccination against influenza is recommended for....adults and children who have immunosuppression (including immunosuppression caused by medications or by human immunodeficiency virus)." ²	
Use in Other Federal Programs:	Centers for Medicare and Medicaid Services EHR Incentive Program measure (number 0041). See https://www.cms.gov/regulations-and-guidance/legislation/ehrincentiveprograms/ecqm_library.html	
References/ Notes:	¹ The HIV/AIDS Bureau did not develop this measure. The American Medical Association-convened Physician Consortium for Performance Improvement (AMA-PCPI) developed this measure. More details available at: https://www.cms.gov/regulations-and-guidance/legislation/ehrincentiveprograms/ecqm_library.html	



HIV/AIDS Bureau Performance Measures



	<p>²Centers for Disease Control and Prevention. Prevention and Control of Influenza: Recommendations from the Advisory committee on Immunization Practices (ACIP). MMWR 2007; 56(RR#6)[1-60]. Available at: http://www.cdc.gov/mmwr/PDF/rr/rr5606.pdf. Section accessed August 14, 2013. Page 2.</p>
--	---



HIV/AIDS Bureau Performance Measures



Performance Measure:	Lipid Screening	National Quality Forum #: None
Percentage of patients, regardless of age, with a diagnosis of HIV who were prescribed HIV antiretroviral therapy and who had a fasting lipid ¹ panel during the measurement year		
Numerator:	Number of patients who had a fasting lipid panel in the measurement year	
Denominator:	Number of patients, regardless of age, who are prescribed HIV antiretroviral therapy and who had a medical visit with a provider with prescribing privileges ² at least once in the measurement year	
Patient Exclusions:	None	
Data Elements:	<ol style="list-style-type: none"> 1. Does the patient, regardless of age, have a diagnosis of HIV? (Y/N) <ol style="list-style-type: none"> a. If yes, did the patient have at least one medical visit during the measurement year? (Y/N) <ol style="list-style-type: none"> i. If yes, was the client prescribed HIV antiretroviral therapy during the measurement year?(Y/N) <ol style="list-style-type: none"> 1. If yes, did the patient have a fasting lipid panel during the measurement year? (Y/N) 	
Comparison Data:	<p>National HIVQUAL:</p> <p>2007: Among patients who were on HAART during the review period, percentage who had cholesterol & triglycerides checked during the review period</p> <p>2009: Percentage of patients who had a lipid profile (cholesterol & triglycerides checked) during the review period</p> <p>2011: Percentage of patients for whom a lipid screening was performed during the review period. At a minimum, lipid screening should include determination of cholesterol, high-density lipoprotein (HDL) and triglyceride levels (https://www.ehivqual.org/ and http://nationalqualitycenter.org/). The National HIVQUAL reported the median as 91% in 2007, 86% in 2009, and 83% in 2011.</p>	
U.S. Department of Health & Human Services Guidelines:	<p><u>Adult guidelines:</u>³ Fasting lipid profile should be monitored for patients at the following points: entry into care; follow up before ART; ART initiation or modification; 2 weeks post ART initiation; every 6 months; every 12 months; and if clinically indicated.</p> <p><u>Pediatric guidelines:</u>⁴ Lipid Panel should be monitored for children during the following points of care: entry into care; ART initiation; and every 6-12 months.</p>	
Use in Other Federal Programs:	None	
References/ Notes:	<p>¹A fasting lipid panel consists of fasting cholesterol, HDL, calculated LDL and triglycerides.</p> <p>²A “provider with prescribing privileges” is a health care professional who is certified in their jurisdiction to prescribe ARV therapy.</p> <p>³Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the use of</p>	



HIV/AIDS Bureau Performance Measures



	<p>antiretroviral agents in HIV-1-infected adults and adolescents. Department of Health and Human Services. Available at http://aidsinfo.nih.gov/ContentFiles/AdultandAdolescentGL.pdf. Section accessed October 14 2014. C-3.</p> <p>⁴Panel on Antiretroviral Therapy and Medical Management of HIV-Infected Children. Guidelines for the Use of Antiretroviral Agents in Pediatric HIV Infection. Available at https://aidsinfo.nih.gov/guidelines. Section accessed October 14, 2014. H-1-3.</p>
--	--



HIV/AIDS Bureau Performance Measures



Performance Measure:	Tuberculosis Screening¹	National Quality Forum #: 408
Percentage of patients aged 3 months and older with a diagnosis of HIV/AIDS, for whom there was documentation that a tuberculosis (TB) screening test was performed and results interpreted (for tuberculin skin tests) at least once since the diagnosis of HIV infection		
Numerator:	<p>Patients for whom there was documentation that a tuberculosis (TB) screening test was performed and results interpreted (for tuberculin skin tests) at least once since the diagnosis of HIV infection.</p> <p>NOTE: Results from the tuberculin skin test must be interpreted by a health care professional.</p>	
Denominator:	All patients aged 3 months and older with a diagnosis of HIV/AIDS, who had at least two visits during the measurement year, with at least 90 days in between each visit.	
Patient Exclusions:	1. Documentation of Medical Reason for not performing a tuberculosis (TB) screening test (e.g., patients with a history of positive PPD or treatment for TB)	
Data Elements:	<p>1. Does the patient, aged three months and older, have a diagnosis of HIV/AIDS? (Y/N)</p> <p>a. If yes, did the patient have at least two medical visits during the measurement year, with at least 90 days in between each visit? (Y/N)</p> <p>i. If yes, has the patient had tuberculosis (TB) screening test performed and results interpreted (for tuberculin skin tests) at least once since the diagnosis of HIV infection? (Y/N)</p>	
Comparison Data:	<p>National HIVQUAL: 2006 & 2007: Percentage of patients without prior positive test or TB treatment who received a TB test with documented result during the past 24 months 2011: Percentage of patient for whom an LTBI screening was performed and the results were read either during the review period or the twelve months preceding the start of the review period (https://www.ehivqual.org/ and http://nationalqualitycenter.org/). The National HIVQUAL reported the median percentage as: 75% in 2007, 74% in 2009, 73% in 2011.</p>	
U.S. Department of Health & Human Services Guidelines:	<p><u>Adult guidelines:</u>² “Testing for LTBI at the time of HIV diagnosis should be routine, regardless of an individual’s epidemiological risk of TB exposure. Individuals with negative diagnostic tests for LTBI who have advanced HIV infection (CD4 cell count <200 cells/mm³) and no indications for initiating empiric LTBI treatment should be retested for LTBI once they start ART and attain a CD4 count ≥200 cells/mm³. Annual testing for LTBI is recommended only for HIV-infected patients who are at high risk of repeated or ongoing exposure to those with active TB.”</p> <p><u>Pediatric guideline:</u>³ The cornerstone of diagnostic methods for latent TB infection (LTBI) is the tuberculin skin test (TST), administered by the Mantoux method. Because children with HIV infection are at high risk for TB, annual testing of this population is recommended to diagnose LTBI (AIII).</p>	



HIV/AIDS Bureau Performance Measures



Use in Other Federal Programs:	None
References/Notes:	<p>¹The HIV/AIDS Bureau did not develop this measure. The National Committee on Quality Assurance developed this measure. Measure details available at: http://www.qualityforum.org/News_And_Resources/Endorsement_Summaries/Endorsement_Summaries.aspx</p> <p>²Panel on Opportunistic Infections in HIV-Infected Adults and Adolescents. Guidelines for the prevention and treatment of opportunistic infections in HIV-infected adults and adolescents: recommendations from the Centers for Disease Control and Prevention, the National Institutes of Health, and the HIV Medicine Association of the Infectious Diseases Society of America. Available at https://aidsinfo.nih.gov/guidelines. Section accessed October 14, 2014. F-2.</p> <p>³Centers for Disease Control and Prevention. Guidelines for the Prevention and Treatment of Opportunistic Infections Among HIV-Exposed and HIV-Infected Children. MMWR 2009;58(No. RR-11). https://aidsinfo.nih.gov/guidelines. Section accessed August 24, 2013. Page 20.</p>

Accessibility: If you need an alternative means of access to any information above please contact us at comments@hrsa.gov. Please let us know the nature of your accessibility problem and the Web address of the requested information.